

Predikaatloogika sekvensiarvutus

- Predikaatloogika sekvensiarvutuses lisanduvad lauseloogika sekvensiarvutusele järgmised reeglid kvantorite kohta:

$$\frac{\frac{\Gamma \rightarrow A[y/x], \Delta}{\Gamma \rightarrow \forall x A, \Delta} \forall \mathcal{R}^*}{\frac{\Gamma \rightarrow A[t/x], (\exists x A,) \Delta}{\Gamma \rightarrow \exists x A, \Delta} \exists \mathcal{R}}$$

$$\frac{\frac{\Gamma, (\forall x A,) A[t/x] \rightarrow \Delta}{\Gamma, \forall x A \rightarrow \Delta} \forall \mathcal{L}}{\frac{\Gamma, A[y/x] \rightarrow \Delta}{\Gamma, \exists x, A \rightarrow \Delta} \exists \mathcal{L}^\dagger}$$

* y ei tohi vabalt esineda valemis $\forall x A$ ja valemihulkades Γ, Δ

† y ei tohi vabalt esineda valemis $\exists x A$ ja valemihulkades Γ, Δ

- Näiteid tõestustest:

$$\begin{array}{c}
 \frac{}{p(x', y') \rightarrow p(x', y')} \text{ hyp.} \\
 \frac{}{\forall y p(x', y) \rightarrow p(x', y')} \forall \mathcal{L} \\
 \frac{}{\forall y p(x', y) \rightarrow \exists x p(x, y')} \exists \mathcal{R} \\
 \frac{}{\exists x \forall y p(x, y) \rightarrow \exists x p(x, y')} \exists \mathcal{L}, x' \text{ värske} \\
 \frac{}{\exists x \forall y p(x, y) \rightarrow \forall y \exists x p(x, y)} \forall \mathcal{R}, y' \text{ värske} \\
 \rightarrow \exists x \forall y p(x, y) \supset \forall y \exists x p(x, y) \supset \mathcal{R}
 \end{array}$$

$$\begin{array}{c}
 \frac{}{p(x') \rightarrow p(x')} \text{ hyp.} \\
 \frac{}{p(x'), \neg p(x') \rightarrow} \neg \mathcal{L} \\
 \frac{}{p(x'), \forall x \neg p(x) \rightarrow} \forall \mathcal{L} \\
 \frac{}{\exists x p(x), \forall x \neg p(x) \rightarrow} \exists \mathcal{L}, x' \text{ värske} \\
 \frac{}{\exists x p(x) \rightarrow \neg \forall x \neg p(x)} \neg \mathcal{R} \\
 \rightarrow \exists x p(x) \supset \neg \forall x \neg p(x) \supset \mathcal{R}
 \end{array}$$

$$\begin{array}{c}
\frac{\overline{p(f(x'), y') \rightarrow p(f(x'), y')}}{\overline{p(f(x'), y') \rightarrow \exists y p(f(x'), y)}} \text{hyp.} \\
\frac{\overline{p(f(x'), y') \rightarrow \exists y p(f(x'), y)}}{\overline{p(f(x'), y') \rightarrow \exists x \exists y p(x, y)}} \exists \mathcal{R} \\
\frac{\overline{p(f(x'), y') \rightarrow \exists x \exists y p(x, y)}}{\overline{p(f(x'), y') \vee p(y', g(x')) \rightarrow \exists x \exists y p(x, y)}} \exists \mathcal{R} \\
\frac{\overline{p(f(x'), y') \vee p(y', g(x')) \rightarrow \exists x \exists y p(x, y)}}{\overline{\exists y (p(f(x'), y) \vee p(y, g(x')))) \rightarrow \exists x \exists y p(x, y)}} \vee \mathcal{L} \\
\frac{\overline{\exists y (p(f(x'), y) \vee p(y, g(x')))) \rightarrow \exists x \exists y p(x, y)}}{\overline{\exists x \exists y (p(f(x), y) \vee p(y, g(x))) \rightarrow \exists x \exists y p(x, y)}} \exists \mathcal{L}, y' \text{ v\u00e4rske} \\
\frac{\overline{\exists x \exists y (p(f(x), y) \vee p(y, g(x))) \rightarrow \exists x \exists y p(x, y)}}{\overline{\rightarrow \exists x \exists y (p(f(x), y) \vee p(y, g(x))) \supset \exists x \exists y p(x, y)}} \exists \mathcal{L}, x' \text{ v\u00e4rske} \\
\rightarrow \exists x \exists y (p(f(x), y) \vee p(y, g(x))) \supset \exists x \exists y p(x, y) \supset \mathcal{R}
\end{array}$$