

Generating Functions

ITT9131 Konkreetne Matemaatika

Chapter Seven

Domino Theory and Change

Basic Maneuvers

Solving Recurrences

Special Generating Functions

Convolutions

Exponential Generating Functions

Dirichlet Generating Functions



1 Convolutions

- Fibonacci convolution
- m -fold convolution
- Catalan numbers

2 Exponential generating functions



Next section

1 Convolutions

- Fibonacci convolution
- m -fold convolution
- Catalan numbers

2 Exponential generating functions



Convolutions

- Given two sequences:

$$\langle f_0, f_1, f_2, \dots \rangle = \langle f_n \rangle \text{ and } \langle g_0, g_1, g_2, \dots \rangle = \langle g_n \rangle$$

The **convolution** of $\langle f_n \rangle$ and $\langle g_n \rangle$ is the sequence

$$\langle f_0 g_0, f_0 g_1 + f_1 g_0, f_0 g_2 + f_1 g_1 + f_2 g_0, \dots \rangle = \left\langle \sum_k f_k g_{n-k} \right\rangle = \left\langle \sum_{k+\ell=n} f_k g_\ell \right\rangle.$$

- If $F(z)$ and $G(z)$ are generating functions on the sequences $\langle f_n \rangle$ and $\langle g_n \rangle$, then their convolution has the generating function $F(z) \cdot G(z)$.
- Three or more sequences can be convolved analogously, for example:

$$\langle f_n \rangle \langle g_n \rangle \langle h_n \rangle = \left\langle \sum_{j+k+\ell=n} f_j g_k h_\ell \right\rangle$$

and the generating function of this three-fold convolution is the product $F(z) \cdot G(z) \cdot H(z)$.



Convolutions

- Given two sequences:

$$\langle f_0, f_1, f_2, \dots \rangle = \langle f_n \rangle \text{ and } \langle g_0, g_1, g_2, \dots \rangle = \langle g_n \rangle$$

The **convolution** of $\langle f_n \rangle$ and $\langle g_n \rangle$ is the sequence

$$\langle f_0 g_0, f_0 g_1 + f_1 g_0, f_0 g_2 + f_1 g_1 + f_2 g_0, \dots \rangle = \left\langle \sum_k f_k g_{n-k} \right\rangle = \left\langle \sum_{k+\ell=n} f_k g_\ell \right\rangle.$$

- If $F(z)$ and $G(z)$ are generating functions on the sequences $\langle f_n \rangle$ and $\langle g_n \rangle$, then their convolution has the generating function $F(z) \cdot G(z)$.
- Three or more sequences can be convolved analogously, for example:

$$\langle f_n \rangle \langle g_n \rangle \langle h_n \rangle = \left\langle \sum_{j+k+\ell=n} f_j g_k h_\ell \right\rangle$$

and the generating function of this three-fold convolution is the product $F(z) \cdot G(z) \cdot H(z)$.



Convolutions

- Given two sequences:

$$\langle f_0, f_1, f_2, \dots \rangle = \langle f_n \rangle \text{ and } \langle g_0, g_1, g_2, \dots \rangle = \langle g_n \rangle$$

The **convolution** of $\langle f_n \rangle$ and $\langle g_n \rangle$ is the sequence

$$\langle f_0 g_0, f_0 g_1 + f_1 g_0, f_0 g_2 + f_1 g_1 + f_2 g_0, \dots \rangle = \left\langle \sum_k f_k g_{n-k} \right\rangle = \left\langle \sum_{k+l=n} f_k g_l \right\rangle.$$

- If $F(z)$ and $G(z)$ are generating functions on the sequences $\langle f_n \rangle$ and $\langle g_n \rangle$, then their convolution has the generating function $F(z) \cdot G(z)$.
- Three or more sequences can be convolved analogously, for example:

$$\langle f_n \rangle \langle g_n \rangle \langle h_n \rangle = \left\langle \sum_{j+k+l=n} f_j g_k h_l \right\rangle$$

and the generating function of this three-fold convolution is the product $F(z) \cdot G(z) \cdot H(z)$.



Next subsection

1 Convolutions

- Fibonacci convolution
- m -fold convolution
- Catalan numbers

2 Exponential generating functions



Fibonacci convolution

To compute $\sum_k f_k f_{n-k}$ use Fibonacci generating function (in the form given by Theorem 1 and considering that $\sum(n+1)z^n = \frac{1}{(1-z)^2}$):

$$\begin{aligned}F^2(z) &= \left(\frac{1}{\sqrt{5}} \left(\frac{1}{1-\Phi z} - \frac{1}{1-\widehat{\Phi} z} \right) \right)^2 \\&= \frac{1}{5} \left(\frac{1}{(1-\Phi z)^2} - \frac{2}{(1-\Phi z)(1-\widehat{\Phi} z)} + \frac{1}{(1-\widehat{\Phi} z)^2} \right) \\&= \frac{1}{5} \sum_{n \geq 0} (n+1)\Phi^n z^n - \frac{2}{5} \sum_{n \geq 0} f_{n+1} z^n + \frac{1}{5} \sum_{n \geq 0} (n+1)\widehat{\Phi}^n z^n \\&= \frac{1}{5} \sum_{n \geq 0} (n+1)(\Phi^n + \widehat{\Phi}^n) z^n - \frac{2}{5} \sum_{n \geq 0} f_{n+1} z^n \\&= \frac{1}{5} \sum_{n \geq 0} (n+1)(2f_{n+1} - f_n) z^n - \frac{2}{5} \sum_{n \geq 0} f_{n+1} z^n \\&= \frac{1}{5} \sum_{n \geq 0} (2nf_{n+1} - (n+1)f_n) z^n\end{aligned}$$

Hence

$$\sum_k f_k f_{n-k} = \frac{2nf_{n+1} - (n+1)f_n}{5}$$



Fibonacci convolution (2)

On the previous slide the following was used:

Property

For any $n \geq 0$: $\Phi^n + \widehat{\Phi}^n = 2f_{n+1} - f_n$

Proof

The equalities $\sum_n \Phi^n z^n = \frac{1}{1-\Phi z}$, $\sum_n \widehat{\Phi}^n z^n = \frac{1}{1-\widehat{\Phi} z}$, and $\Phi + \widehat{\Phi} = 1$ are used in the following derivation:

$$\begin{aligned}\sum_n (\Phi^n + \widehat{\Phi}^n) z^n &= \frac{1}{1-\Phi z} + \frac{1}{1-\widehat{\Phi} z} = \frac{1-\widehat{\Phi} z + 1-\Phi z}{(1-\Phi z)(1-\widehat{\Phi} z)} = \\ &= \frac{2-z}{1-z-z^2} = \frac{2}{z} \cdot \frac{z}{1-z-z^2} - \frac{z}{1-z-z^2} = \\ &= \frac{2}{z} \sum_n f_n z^n - \sum_n f_n z^n = 2 \sum_n f_n z^{n-1} - \sum_n f_n z^n = \\ &= 2 \sum_n f_{n+1} z^n - \sum_n f_n z^n = \\ &= \sum_n (2f_{n+1} - f_n) z^n\end{aligned}$$



Fibonacci convolution (2)

On the previous slide the following was used:

Property

For any $n \geq 0$: $\Phi^n + \widehat{\Phi}^n = 2f_{n+1} - f_n$

Proof (alternative)

We know from Exercise 6.28 that

$$\Phi^n + \widehat{\Phi}^n = L_n = f_{n+1} + f_{n-1},$$

with the convention $f_{-1} = 1$, is the n th Lucas number, which is the solution to the recurrence:

$$\begin{aligned} L_0 &= 2; & L_1 &= 1; \\ L_n &= L_{n-1} + L_{n-2} & \forall n &\geq 2. \end{aligned}$$

By writing the recurrence relation for Fibonacci numbers in the form $f_{n-1} = f_{n+1} - f_n$ (which, incidentally, yields $f_{-1} = 1$), we get precisely $L_n = 2f_{n+1} - f_n$.

Q.E.D.



Next subsection

1 Convolutions


- Fibonacci convolution
- m -fold convolution
- Catalan numbers

2 Exponential generating functions





Spanning trees for fan

Example: the fan of order 5:




Spanning trees:





Spanning trees for fan

Example: the fan of order 5:




Spanning trees:



$f_3 = 8$

Spanning trees for fan (2)




How many spanning trees can we make?

- We need to connect 0 to each of the four blocks:
 - two ways to join 0 with $\{9, 10\}$,
 - one way to join 0 with $\{8\}$,
 - four ways to join 0 with $\{4, 5, 6, 7\}$,
 - three ways to join 0 with $\{1, 2, 3\}$,
- There is altogether $2 \cdot 1 \cdot 4 \cdot 3 = 24$ ways for that.

In general:

$$s_n = \sum_{m \geq 0} \sum_{\substack{k_1 + k_2 + \dots + k_m = n \\ k_1, k_2, \dots, k_m > 0}} k_1 k_2 \dots k_m$$

Spanning trees for fan (2)



How many spanning trees can we make?

- We need to connect 0 to each of the four blocks:
 - two ways to join 0 with {9,10},
 - one way to join 0 with {8},
 - four ways to join 0 with {4,5,6,7},
 - three ways to join 0 with {1,2,3},
- There is altogether $2 \cdot 1 \cdot 4 \cdot 3 = 24$ ways for that.

In general:


$$s_n = \sum_{m>0} \sum_{\substack{k_1+k_2+\dots+k_m=n \\ k_1, k_2, \dots, k_m > 0}} k_1 k_2 \dots k_m$$

For example

$$f_4 = 4 + \underbrace{3 \cdot 1 + 2 \cdot 2 + 1 \cdot 3}_{= 10} + \underbrace{2 \cdot 1 \cdot 1 + 1 \cdot 2 \cdot 1 + 1 \cdot 1 \cdot 2}_{= 6} + 1 \cdot 1 \cdot 1 \cdot 1 = 21$$



Spanning trees for fan (2)



How many spanning trees can we make?

- We need to connect 0 to each of the four blocks:
 - two ways to join 0 with $\{9, 10\}$,
 - one way to join 0 with $\{8\}$,
 - four ways to join 0 with $\{4, 5, 6, 7\}$,
 - three ways to join 0 with $\{1, 2, 3\}$,
- There is altogether $2 \cdot 1 \cdot 4 \cdot 3 = 24$ ways for that.

In general:

$$s_n = \sum_{m>0} \sum_{\substack{k_1 + k_2 + \dots + k_m = n \\ k_1, k_2, \dots, k_m > 0}} k_1 k_2 \dots k_m$$

This is the sum of m -fold convolutions of the sequence $\langle 0, 1, 2, 3, \dots \rangle$.



Spanning trees for fan (3)

Generating function for the number of spanning trees:

- The sequence $\langle 0, 1, 2, 3, \dots \rangle$ has the generating function

$$G(z) = \frac{z}{(1-z)^2}.$$

- Hence the generating function for $\langle f_n \rangle$ is

$$\begin{aligned} S(z) &= G(z) + G^2(z) + G^3(z) + \dots = \frac{G(z)}{1-G(z)} \\ &= \frac{z}{(1-z)^2 \left(1 - \frac{z}{(1-z)^2}\right)} \\ &= \frac{z}{(1-z)^2 - z} \\ &= \frac{z}{1-3z+z^2}. \end{aligned}$$



Spanning trees for fan (3)

Generating function for the number of spanning trees:

- The sequence $\langle 0, 1, 2, 3, \dots \rangle$ has the generating function

$$G(z) = \frac{z}{(1-z)^2}.$$

- Hence the generating function for $\langle f_n \rangle$ is

$$\begin{aligned} S(z) &= G(z) + G^2(z) + G^3(z) + \dots = \frac{G(z)}{1-G(z)} \\ &= \frac{z}{(1-z)^2 \left(1 - \frac{z}{(1-z)^2}\right)} \\ &= \frac{z}{(1-z)^2 - z} \\ &= \frac{z}{1-3z+z^2}. \end{aligned}$$

Consequently $s_n = f_{2n}$



Next subsection

1 Convolutions

- Fibonacci convolution
- m -fold convolution
- Catalan numbers

2 Exponential generating functions



Dyck language

Definition

The Dyck language is the language consisting of balanced strings of parentheses '[' and ']'.

Another definition

If $X = \{x, \bar{x}\}$ is the alphabet, then the **Dyck language** is the subset \mathcal{D} of words u of X^* which satisfy

- $|u|_x = |u|_{\bar{x}}$, where $|u|_x$ is the number of letters x in the word u , and
- if u is factored as vw , where v and w are words of X^* , then $|v|_x \geq |v|_{\bar{x}}$.

	0 pairs	1 pair	2 pairs	3 pairs
Elements	\emptyset	[]	[[]] [] []	[[[]]] [[] [] [] [[]] [[] [] [] [] []
	\emptyset	AB	AABB ABAB	AAABBB AABBBAB ABAABB AABABB ABABAB
No of words	1	1	2	5



Dyck language (2)

- Let C_n be the number of words in the Dyck language \mathcal{D} having exactly n pairs of parentheses.
- If $u = vw$ for $u \in \mathcal{D}$, then the prefix $v \in \mathcal{D}$ iff the suffix $w \in \mathcal{D}$
- Then every word $u \in \mathcal{D}$ of length ≥ 2 has a unique writing $u = [v]w$ such that $v, w \in \mathcal{D}$ (possibly empty) but $[p \notin \mathcal{D}$ for every prefix p of u (including u itself).
- Hence, for any $n > 0$

$$C_n = C_0 C_{n-1} + C_1 C_{n-2} + \cdots + C_{n-1} C_0$$

- The number series $\langle C_n \rangle$ is called **Catalan numbers**, from the Belgian mathematician Eugène Catalan.
Let us derive the closed formula for C_n in the following slides.



Catalan numbers

Step 1 The recurrent equation of Catalan numbers for all integers

$$C_n = \sum_k C_k C_{n-1-k} + [n=0].$$

Step 2 Write down $C(z) = \sum_n C_n z^n$:

$$\begin{aligned} C(z) &= \sum_n C_n z^n = \sum_{k,n} C_k C_{n-1-k} z^n + \sum_n [n=0] z^n \\ &= \sum_k C_k z^k z \sum_n C_{n-1-k} z^{n-1-k} + 1 \\ &= \sum_k C_k z^k z \sum_n C_n z^n + 1 \\ &= zC^2(z) + 1 \end{aligned}$$



Catalan numbers

Step 1 The recurrent equation of Catalan numbers for all integers

$$C_n = \sum_k C_k C_{n-1-k} + [n = 0].$$

Step 2 Write down $C(z) = \sum_n C_n z^n$:

$$\begin{aligned} C(z) &= \sum_n C_n z^n = \sum_{k,n} C_k C_{n-1-k} z^n + \sum_n [n = 0] z^n \\ &= \sum_k C_k z^k z \sum_n C_{n-1-k} z^{n-1-k} + 1 \\ &= \sum_k C_k z^k z \sum_n C_n z^n + 1 \\ &= zC^2(z) + 1 \end{aligned}$$



Catalan numbers (2)

Step 3 Solving the quadratic equation $zC^2(z) - C(z) + 1 = 0$ for $C(z)$ yields directly:

$$C(z) = \frac{1 \pm \sqrt{1-4z}}{2z}.$$

(Solution with "+" isn't proper as it leads to $C_0 = C(0) = \infty$.)

Step 4 From the binomial theorem we get:

$$\sqrt{1-4z} = \sum_{k \geq 0} \binom{1/2}{k} (-4z)^k = 1 + \sum_{k \geq 1} \frac{1}{2k} \binom{-1/2}{k-1} (-4z)^k$$

- Using the equality for binomials $\binom{-1/2}{n} = (-1/4)^n \binom{2n}{n}$ we finally get

$$\begin{aligned} C(z) &= \frac{1 - \sqrt{1-4z}}{2z} = \sum_{k \geq 1} \frac{1}{k} \binom{-1/2}{k-1} (-4z)^{k-1} \\ &= \sum_{n \geq 0} \binom{-1/2}{n} \frac{(-4z)^n}{n+1} \\ &= \sum_{n \geq 0} \binom{2n}{n} \frac{z^n}{n+1} \end{aligned}$$



Catalan numbers (2)

Step 3 Solving the quadratic equation $zC^2(z) - C(z) + 1 = 0$ for $C(z)$ yields directly:

$$C(z) = \frac{1 \pm \sqrt{1-4z}}{2z}.$$

(Solution with "+" isn't proper as it leads to $C_0 = C(0) = \infty$.)

Step 4 From the binomial theorem we get:

$$\sqrt{1-4z} = \sum_{k \geq 0} \binom{1/2}{k} (-4z)^k = 1 + \sum_{k \geq 1} \frac{1}{2k} \binom{-1/2}{k-1} (-4z)^k$$

- Using the equality for binomials $\binom{-1/2}{n} = (-1/4)^n \binom{2n}{n}$ we finally get

$$\begin{aligned} C(z) &= \frac{1 - \sqrt{1-4z}}{2z} = \sum_{k \geq 1} \frac{1}{k} \binom{-1/2}{k-1} (-4z)^{k-1} \\ &= \sum_{n \geq 0} \binom{-1/2}{n} \frac{(-4z)^n}{n+1} \\ &= \sum_{n \geq 0} \binom{2n}{n} \frac{z^n}{n+1} \end{aligned}$$



Proof that $\binom{-1/2}{n} = (-1/4)^n \binom{2n}{n}$

We prove a bit more: for every $r \in \mathbb{R}$ and $k \geq 0$,

$$r^k \cdot \left(r - \frac{1}{2}\right)^k = \frac{(2r)^{2k}}{2^{2k}}$$



Proof that $\binom{-1/2}{n} = (-1/4)^n \binom{2n}{n}$

We prove a bit more: for every $r \in \mathbb{R}$ and $k \geq 0$,

$$r^k \cdot \left(r - \frac{1}{2}\right)^k = \frac{(2r)^{2k}}{2^{2k}}$$

Indeed,

$$\begin{aligned} r^k \cdot \left(r - \frac{1}{2}\right)^k &= r \cdot \left(r - \frac{1}{2}\right) \cdot (r-1) \cdot \left(r - \frac{3}{2}\right) \cdots (r-k-1) \cdot \left(r - \frac{1}{2} - k + 1\right) \\ &= \frac{2r}{2} \cdot \frac{2r-1}{2} \cdot \frac{2r-2}{2} \cdot \frac{2r-3}{2} \cdots \frac{2r-2k-2}{2} \cdot \frac{2r-2k+1}{2} \\ &= \frac{(2r)^{2k}}{2^{2k}} \end{aligned}$$



Proof that $\binom{-1/2}{n} = (-1/4)^n \binom{2n}{n}$

We prove a bit more: for every $r \in \mathbb{R}$ and $k \geq 0$,

$$r^k \cdot \left(r - \frac{1}{2}\right)^k = \frac{(2r)^{2k}}{2^{2k}}$$

Indeed,

$$\begin{aligned} r^k \cdot \left(r - \frac{1}{2}\right)^k &= r \cdot \left(r - \frac{1}{2}\right) \cdot (r-1) \cdot \left(r - \frac{3}{2}\right) \cdots (r-k-1) \cdot \left(r - \frac{1}{2} - k + 1\right) \\ &= \frac{2r}{2} \cdot \frac{2r-1}{2} \cdot \frac{2r-2}{2} \cdot \frac{2r-3}{2} \cdots \frac{2r-2k-2}{2} \cdot \frac{2r-2k+1}{2} \\ &= \frac{(2r)^{2k}}{2^{2k}} \end{aligned}$$

Then for $r = k = n$, dividing by $(n!)^2$ and using $n^n = n!$,

$$\binom{n-1/2}{n} = \left(\frac{1}{4}\right)^n \binom{2n}{n} :$$

and as $r^k = (-1)^k (-r)^{\overline{k}} = (-1)^k (-r+k-1)^{\overline{k}}$,

$$\binom{-1/2}{n} = \binom{n - (n-1/2) - 1}{n} = \frac{(-1)^n}{4^n} \binom{2n}{n}$$



Resume Catalan numbers

Formulae for computation

- $C_{n+1} = \frac{2(2n+1)}{n+2} C_n$, with $C_0 = 1$
- $C_n = \frac{1}{n+1} \binom{2n}{n}$
- $C_n = \binom{2n}{n} - \binom{2n}{n-1} = \binom{2n-1}{n} - \binom{2n-1}{n+1}$
- Generating function: $C(z) = \frac{1-\sqrt{1-4z}}{2z}$



Eugène Charles Catalan
(1814–1894)


$$\lim_{n \rightarrow \infty} \frac{C_n}{C_{n-1}} = 4$$

n	0	1	2	3	4	5	6	7	8	9	10
C_n	1	1	2	5	14	42	132	429	1 430	4 862	16 796



Applications of Catalan numbers

Number of complete binary trees with $n+1$ leaves is C_n



The **Dyck language** consists of exactly n characters A and n characters B, and every prefix does not contain more B-s than A-s. For example, there are five words with 6 letters in the Dyck language:

AAABBB AABABB AABBAB ABAABB ABABAB


Corollary

C_n is the number of words of length $2n$ in the Dyck language.



Applications of Catalan numbers

Number of complete binary trees with $n+1$ leaves is C_n



The **Dyck language** consists of exactly n characters A and n characters B, and every prefix does not contain more B-s than A-s. For example, there are five words with 6 letters in the Dyck language:

AAABBB AABABB AABBAB ABAABB ABABAB

Corollary


C_n is the number of words of length $2n$ in the Dyck language.



Applications of Catalan numbers (2)

Monotonic paths


C_n is the number of monotonic paths along the edges of a grid with $n \times n$ square cells, which do not pass above the diagonal. A monotonic path is one which starts in the lower left corner, finishes in the upper right corner, and consists entirely of edges pointing rightwards or upwards.



Applications of Catalan numbers (3)

Polygon triangulation

C_n is the number of different ways a convex polygon with $n+2$ sides can be cut into triangles by connecting vertices with straight lines.



See more applications, for example, on

http://www.absoluteastronomy.com/topics/Catalan_number



Next section

- 1 Convolutions
 - Fibonacci convolution
 - m -fold convolution
 - Catalan numbers
- 2 Exponential generating functions



Exponential generating function

Definition

The **exponential generating function** (briefly, egf) of the sequence $\langle g_n \rangle$ is the function

$$\widehat{G}(z) = \sum_{n \geq 0} \frac{g_n}{n!} z^n,$$

that is, the generating function of the sequence $\langle g_n/n! \rangle$.

For example, $e^z = \sum_{n \geq 0} \frac{z^n}{n!}$ is the egf of the constant sequence 1.



Exponential generating function

Definition

The **exponential generating function** (briefly, egf) of the sequence $\langle g_n \rangle$ is the function

$$\widehat{G}(z) = \sum_{n \geq 0} \frac{g_n}{n!} z^n,$$

that is, the generating function of the sequence $\langle g_n/n! \rangle$.

For example, $e^z = \sum_{n \geq 0} \frac{z^n}{n!}$ is the egf of the constant sequence 1.

Why exponential generating functions?

Because $\langle g_n/n! \rangle$ might have a “simpler” generating function than $\langle g_n \rangle$ has.



Exponential generating functions: Basic maneuvers

Let $\hat{F}(z)$ and $\hat{G}(z)$ be the exponential generating functions of $\langle f_n \rangle$ and $\langle g_n \rangle$.

As usual, we put $f_n = g_n = 0$ for every $n < 0$, and undefined $\cdot 0 = 0$.

- $\alpha \hat{F}(z) + \beta \hat{G}(z) = \sum_n \left(\frac{\alpha f_n + \beta g_n}{n!} \right) z^n$
- $\hat{G}(cz) = \sum_n \frac{c^n g_n}{n!} z^n$
- $z \hat{G}(z) = \sum_n \frac{n g_{n-1}}{n!} z^n$
- $\hat{G}'(z) = \sum_n \frac{g_{n+1}}{n!} z^n$
- $\int_0^z \hat{G}(w) dw = \sum_n \frac{g_{n-1}}{n!} z^n$
- $\hat{F}(z) \cdot \hat{G}(z) = \sum_n \frac{1}{n!} \left(\sum_k \binom{n}{k} f_k g_{n-k} \right) z^n$



Exponential generating functions: Basic maneuvers

Let $\hat{F}(z)$ and $\hat{G}(z)$ be the exponential generating functions of $\langle f_n \rangle$ and $\langle g_n \rangle$.

As usual, we put $f_n = g_n = 0$ for every $n < 0$, and undefined $\cdot 0 = 0$.

$$\blacksquare \alpha \hat{F}(z) + \beta \hat{G}(z) = \sum_n \left(\frac{\alpha f_n + \beta g_n}{n!} \right) z^n$$

$$\blacksquare \hat{G}(cz) = \sum_n \frac{c^n g_n}{n!} z^n$$

$$\blacksquare z \hat{G}(z) = \sum_n \frac{n g_{n-1}}{n!} z^n$$

$$\blacksquare \hat{G}'(z) = \sum_n \frac{g_{n+1}}{n!} z^n$$

$$\blacksquare \int_0^z \hat{G}(w) dw = \sum_n \frac{g_{n-1}}{n!} z^n$$

$$\blacksquare \hat{F}(z) \cdot \hat{G}(z) = \sum_n \frac{1}{n!} \left(\sum_k \binom{n}{k} f_k g_{n-k} \right) z^n$$



Exponential generating functions: Basic maneuvers

Let $\widehat{F}(z)$ and $\widehat{G}(z)$ be the exponential generating functions of $\langle f_n \rangle$ and $\langle g_n \rangle$.

As usual, we put $f_n = g_n = 0$ for every $n < 0$, and undefined $\cdot 0 = 0$.

- $\alpha \widehat{F}(z) + \beta \widehat{G}(z) = \sum_n \left(\frac{\alpha f_n + \beta g_n}{n!} \right) z^n$
- $\widehat{G}(cz) = \sum_n \frac{c^n g_n}{n!} z^n$
- $z \widehat{G}(z) = \sum_n \frac{n g_{n-1}}{n!} z^n$
- $\widehat{G}'(z) = \sum_n \frac{g_{n+1}}{n!} z^n$
- $\int_0^z \widehat{G}(w) dw = \sum_n \frac{g_{n-1}}{n!} z^n$
- $\widehat{F}(z) \cdot \widehat{G}(z) = \sum_n \frac{1}{n!} \left(\sum_k \binom{n}{k} f_k g_{n-k} \right) z^n$



Exponential generating functions: Basic maneuvers

Let $\widehat{F}(z)$ and $\widehat{G}(z)$ be the exponential generating functions of $\langle f_n \rangle$ and $\langle g_n \rangle$.

As usual, we put $f_n = g_n = 0$ for every $n < 0$, and undefined $\cdot 0 = 0$.

$$\blacksquare \alpha \widehat{F}(z) + \beta \widehat{G}(z) = \sum_n \left(\frac{\alpha f_n + \beta g_n}{n!} \right) z^n$$

$$\blacksquare \widehat{G}(cz) = \sum_n \frac{c^n g_n}{n!} z^n$$

$$\blacksquare z \widehat{G}(z) = \sum_n \frac{n g_{n-1}}{n!} z^n$$

$$\blacksquare \widehat{G}'(z) = \sum_n \frac{g_{n+1}}{n!} z^n$$

$$\blacksquare \int_0^z \widehat{G}(w) dw = \sum_n \frac{g_{n-1}}{n!} z^n$$

$$\blacksquare \widehat{F}(z) \cdot \widehat{G}(z) = \sum_n \frac{1}{n!} \left(\sum_k \binom{n}{k} f_k g_{n-k} \right) z^n$$



Exponential generating functions: Basic maneuvers

Let $\widehat{F}(z)$ and $\widehat{G}(z)$ be the exponential generating functions of $\langle f_n \rangle$ and $\langle g_n \rangle$.

As usual, we put $f_n = g_n = 0$ for every $n < 0$, and undefined $\cdot 0 = 0$.

- $\alpha \widehat{F}(z) + \beta \widehat{G}(z) = \sum_n \left(\frac{\alpha f_n + \beta g_n}{n!} \right) z^n$

- $\widehat{G}(cz) = \sum_n \frac{c^n g_n}{n!} z^n$

- $z \widehat{G}(z) = \sum_n \frac{n g_{n-1}}{n!} z^n$

- $\widehat{G}'(z) = \sum_n \frac{g_{n+1}}{n!} z^n$

- $\int_0^z \widehat{G}(w) dw = \sum_n \frac{g_{n-1}}{n!} z^n$

- $\widehat{F}(z) \cdot \widehat{G}(z) = \sum_n \frac{1}{n!} \left(\sum_k \binom{n}{k} f_k g_{n-k} \right) z^n$



Exponential generating functions: Basic maneuvers

Let $\widehat{F}(z)$ and $\widehat{G}(z)$ be the exponential generating functions of $\langle f_n \rangle$ and $\langle g_n \rangle$.

As usual, we put $f_n = g_n = 0$ for every $n < 0$, and undefined $\cdot 0 = 0$.

- $\alpha \widehat{F}(z) + \beta \widehat{G}(z) = \sum_n \left(\frac{\alpha f_n + \beta g_n}{n!} \right) z^n$
- $\widehat{G}(cz) = \sum_n \frac{c^n g_n}{n!} z^n$
- $z \widehat{G}(z) = \sum_n \frac{n g_{n-1}}{n!} z^n$
- $\widehat{G}'(z) = \sum_n \frac{g_{n+1}}{n!} z^n$
- $\int_0^z \widehat{G}(w) dw = \sum_n \frac{g_{n-1}}{n!} z^n$
- $\widehat{F}(z) \cdot \widehat{G}(z) = \sum_n \frac{1}{n!} \left(\sum_k \binom{n}{k} f_k g_{n-k} \right) z^n$



Binomial convolution

Definition

The **binomial convolution** of the sequences $\langle f_n \rangle$ and $\langle g_n \rangle$ is the sequence $\langle h_n \rangle$ defined by:

$$h_n = \sum_k \binom{n}{k} f_k g_{n-k}$$



Binomial convolution

Definition

The **binomial convolution** of the sequences $\langle f_n \rangle$ and $\langle g_n \rangle$ is the sequence $\langle h_n \rangle$ defined by:

$$h_n = \sum_k \binom{n}{k} f_k g_{n-k}$$

Examples

- $\langle (a+b)^n \rangle$ is the binomial convolution of $\langle a^n \rangle$ and $\langle b^n \rangle$.
- If $\widehat{F}(z)$ is the egf of $\langle f_n \rangle$ and $\widehat{G}(z)$ is the egf of $\langle g_n \rangle$, then $\widehat{H}(z) = \widehat{F}(z) \cdot \widehat{G}(z)$ is the egf of $\langle h_n \rangle$, because then:

$$\frac{h_n}{n!} = \sum_k \frac{f_k}{k!} \frac{g_{n-k}}{(n-k)!}$$



Bernoulli numbers and exponential generating functions

Recall that the **Bernoulli numbers** are defined by the recurrence:

$$\sum_{k=0}^m \binom{m+1}{k} B_k = [m=0] \quad \forall m \geq 0,$$

which is equivalent to:

$$\sum_n \binom{n}{k} B_k = B_n + [n=1] \quad \forall n \geq 0.$$

The left-hand side is a binomial convolution with the constant sequence 1. Then the egf $\widehat{B}(z)$ of the Bernoulli numbers satisfies

$$\widehat{B}(z) \cdot e^z = \widehat{B}(z) + z :$$

which yields

$$\widehat{B}(z) = \frac{z}{e^z - 1}.$$



Bernoulli numbers and exponential generating functions

Recall that the **Bernoulli numbers** are defined by the recurrence:

$$\sum_{k=0}^m \binom{m+1}{k} B_k = [m=0] \quad \forall m \geq 0,$$

which is equivalent to:

$$\sum_n \binom{n}{k} B_k = B_n + [n=1] \quad \forall n \geq 0.$$

To make a comparison:

$$\sum_{n \geq 0} \frac{B_n}{n!} z^n = \frac{z}{e^z - 1} \quad \text{but} \quad \sum_{n \geq 0} B_n^+ z^n = \frac{1}{z} \frac{d^2}{dz^2} \ln \int_0^\infty t^{z-1} e^{-t} dt$$

where $B_n^+ = B_n \cdot [B_n \geq 0]$.

